

## Microtiter Enhanced Binding Scientific References

Hakala J K, K Öörni, M O Pentikäinen, E Hurt-Camejo, P T Kovanen. 2001. Lipolysis of LDL by human secretory phospholipase A2 induces particle fusion and enhances the retention of LDL to human aortic proteoglycans arterioscler. *Thromb. Vasc. Biol.* 21, 1053-1058.

Matsushita K, R Motani, T Sakurai, N Yamaguchi, T Koga, K Matsuo, S Nagaoka, K Abeyama, I Maruyama and M. Torii. 2000. The role of vascular endothelial growth factor in human dental pulp cells: Induction of chemotaxis, proliferation, and differentiation and activation of the AP-1-dependent signaling pathway. *J Dent Res* 79, 1596-1603.

Olin A I, M Mörgelin, T Sasaki, R Timpl, D Heinegård and A Aspberg. 2001. The proteoglycans aggrecan and versican form networks with fibulin-2 through their lectin domain binding. *J Biol Chem* 276, 1253–1261.

Pentikäinen M O, K Öörni and P T Kovanen. 2000. Lipoprotein lipase (LPL) strongly links native and oxidized low density lipoprotein particles to decorin-coated collagen. Roles for both dimeric and monomeric forms of LPL. *J Biol Chem* 275, 5694–5701.

Sneck M, P T Kovanen and K Öörni. 2005. Decrease in pH strongly enhances binding of native, proteolyzed, lipolyzed and oxidized low density lipoprotein particles to human aortic proteoglycans. *J Biol Chem* 280, 37449–37454.

Öörni K, M Sneck, D Brömme, M O Pentikäinen, K A Lindstedt, M Mäyränpää, H Aitio and P T Kovanen. 2004. Cysteine protease cathepsin F is expressed in human atherosclerotic lesions, is secreted by cultured macrophages, and modifies low density lipoprotein particles in vitro. *J Biol Chem* 279, 34776–34784.